Retrieving the missed particle-antiparticle degrees of freedom of Dirac particles

Shun-Jin Wang,[1, 2] Shan-Gui Zhou,[3, 2, *] and Hans-Christian Pauli[4]
[1]*Department of Physics, Sichuan University, Chengdu 610064, China*
[2]*Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China*
[3]*Institute of Theoretical Physics, Chinese Academy Sciences, Beijing 100080, China*
[4]*Max-Planck-Institut für Kernphysik, 69029 Heidelberg, Germany*

(Dated: February 1, 2008)

The missed particle-antiparticle degrees of freedom are retrieved and the corresponding particle-antiparticle intrinsic space are introduced to study the dynamical symmetry of the Dirac particle. As a result, the particle-antiparticle quantum number appears naturally and the Dirac particle has five quantum numbers instead of four. An anti-symmetry (different from the conventional symmetry) of the Dirac Hamiltonian and a dual symmetry of its eigen functions are explored. The $\hat{\kappa}$ operator of the Dirac equation in central potentials is found to be the analog of the helicity operator of the free particle—the alignment of the spin along the angular momentum.

PACS numbers: 03.65.-w, 03.65.Ca, 03.65.Pm, 11.30.Er

arXiv:hep-th/0501250v1 31 Jan 2005

*Introduction.*The eigen solutions of the Dirac equation for a free particle and/or a particle in central potentials are the most basic elements of the relativistic quantum mechanics [1] and quantum field theory [2]. Although the particle and antiparticle states of a Dirac particle are conventionally considered as two states of the same Fermion, and the anti-commutators and the charge conjugate transformation (the $\mathcal{C}$ operator) for the field operators are introduced, the particle-antiparticle quantum number is still missing and the $\mathcal{C}$ transformation just belongs to a discrete two-element group. This unpleasant situation occurs due to the veiled reason that the particle-antiparticle degrees of freedom are hidden in the γ matrices of the Dirac equation. In this Letter we shall show that the γ matrices can be split into products of two parts which correspond to the spin operators and the particle-antiparticle operators respectively. The dynamical degrees of freedom (DoF) for a Dirac particle thus become complete and include those in coordinate space, spin space, and particle-antiparticle space. As a result, the Dirac particle has five quantum numbers instead of four. The splitting of γ matrices and the introduction of particle-antiparticle operators are necessary for a correct description of interactions and may have profound meaning.

The dynamical degrees of freedom for free Dirac particles. The Hamiltonian of a free Dirac particle reads

$$\widehat{H} = c\boldsymbol{\alpha}\cdot\boldsymbol{p} + mc^2\beta. \tag{1}$$

Introducing the particle-antiparticle operators $\boldsymbol{\tau}$ as follows

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\ \tau_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix},\ \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \tag{2}$$

the matrices $\boldsymbol{\alpha}$ and β (or γ_μ) can be written as direct products of the spin operator $\boldsymbol{\sigma}$ and the particle-antiparticle operator $\boldsymbol{\tau}$

$$\boldsymbol{\alpha} = \boldsymbol{\sigma}\otimes\tau_1,\ \beta = \mathbb{I}_\sigma\otimes\tau_3. \tag{3}$$

Here the convention is used that the spin operators $(\mathbb{I}_\sigma, \sigma_i)$ should be inserted into the particle-antiparticle operators $(\mathbb{I}_\tau, \tau_i)$ as their operator-valued elements. It is found that the algebra $(\mathbb{I}_\sigma, \sigma_i)\otimes(\mathbb{I}_\tau, \tau_i)$ is equivalent to the Clifford algebra or Dirac ring which is indispensible for a proper description of interations [3, 4] and also justifies the above splitting scheme. Hence the Hamiltonian can be rewritten as

$$\widehat{H} = c\boldsymbol{p}\cdot\boldsymbol{\sigma}\otimes\tau_1 + mc^2\mathbb{I}_\sigma\otimes\tau_3. \tag{4}$$

Now it is evident that a free Dirac particle has three types of DoF's: the DoF's in coordinate space $\boldsymbol{r}$, spin space s, and particle-antiparticle space τ. Equation (4) clearly shows that the Dirac Hamiltonian possesses the dynamical group $\mathrm{SU}_\sigma(2)\otimes\mathrm{SU}_\tau(2)$. Thus a free Dirac particle should have five mutually commutative invariant operators and five quantum numbers (instead of four) which uniquely specify its quantum states. The five operators are: (i) three components of the conserved momentum $\boldsymbol{p}$ ($[\boldsymbol{p}, \widehat{H}] = 0$); (ii) since the momentum is conserved, the helicity can be defined as $\sigma_\parallel = (\boldsymbol{p}\cdot\boldsymbol{\sigma})/p$ which commutes with the Hamiltonian; (iii) finally, the particle-antiparticle operator reads

$$\widetilde{\tau}_3 = \frac{1}{E}(\mu\ cp\tau_1 + mc^2\tau_3) = \frac{1}{E}\begin{pmatrix} mc^2 & \mu\ cp \\ \mu\ cp & -mc^2 \end{pmatrix}, \tag{5}$$

where μ is the eigen value of $\sigma_\parallel$, and the commutation relation $[\widetilde{\tau}_3, \widehat{H}] = 0$ holds too.

The eigen wave function for a free Dirac particle is a product of three separable parts

$$\widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \tfrac{1}{(2\pi\hbar)^{3/2}}e^{i\boldsymbol{p}\cdot\boldsymbol{r}}\widetilde{U}(\mu)\widetilde{V}(\nu) \tag{6}$$

where the spin wave function $\widetilde{U}(\mu)$ is the eigen function of $\sigma_\parallel$

$$\sigma_\parallel\widetilde{U}(\mu) = \mu\widetilde{U}(\mu),\ \text{with}\ \mu = \pm 1. \tag{7}$$

If $\sigma_\parallel$ is diagonal, *i.e.*, $\sigma_\parallel = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, then we have

$$\widetilde{U}(+1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \ \widetilde{U}(-1) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{8}$$

In the above helicity representation, the eigen functions of $\widetilde{\tau}_3$ with the eigen values $+1$ and -1 corresponding to particle and antiparticle respectively, read

$$\widetilde{V}(+1) = \sqrt{\tfrac{E+mc^2}{2E}} \begin{pmatrix} 1 \\ \frac{\mu cp}{E+mc^2} \end{pmatrix}, \tag{9}$$

$$\widetilde{V}(-1) = i\sqrt{\tfrac{E+mc^2}{2E}} \begin{pmatrix} \frac{-\mu cp}{E+mc^2} \\ 1 \end{pmatrix}, \tag{10}$$

where $E = \sqrt{m^2c^4 + c^2p^2}$. Inserting Eqs. (8, 9, 10) into Eq. (6), we recover the standard solutions for a free Dirac particle.

In the above representation, the 4×4 helicity operator is also diagonal,

$$\widetilde{\Sigma}_\parallel = \mathbb{I}_\tau \otimes \sigma_z, \quad \widetilde{\Sigma}_\parallel \widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \mu \widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau). \tag{11}$$

and the particle-antiparticle operator $\widetilde{\tau}_3$ becomes a 4×4 matrix

$$\widetilde{T}_3 \ = \ \frac{1}{E} \begin{pmatrix} mc^2 & cp\sigma_z \\ cp\sigma_z & -mc^2 \end{pmatrix}, \tag{12}$$

with

$$\widetilde{T}_3 \widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \nu \widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau). \tag{13}$$

It turns out that the Hamiltonian is proportional to the particle-antiparticle operator,

$$\widetilde{H} = E\widetilde{T}_3, \quad \widetilde{H}\widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \nu E \widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau), \tag{14}$$

which tells us that the negative energies of the antiparticles are in fact from the negative antiparticle quantum number -1.

We find that a particle state is transformed to an antiparticle state with the helicity intact by the following unitary transformation,

$$\widetilde{C} = \begin{pmatrix} 0 & i\mathbb{I} \\ -i\mathbb{I} & 0 \end{pmatrix} = \widetilde{C}^\dagger = \widetilde{C}^{-1} \tag{15}$$

$$\widetilde{C}\widetilde{\Psi}_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \widetilde{\Psi}_{p\mu-\nu}(\boldsymbol{r}, s, \tau), \tag{16}$$

and *vice versa*. Thus we can call $\widetilde{C}$ the particle-antiparticle transformation. The Hamiltonian $\widetilde{H}$ is *antisymmetric* under the particle-antiparticle transformation,

$$\widetilde{C}\widetilde{H}\widetilde{C}^{-1} = -\widetilde{H}, \ \textit{i.e.}, \ \{\widetilde{H}, \widetilde{C}\} = 0. \tag{17}$$

By virtue of the unitary transformation,

$$\widehat{u} = \mathbb{I}_\tau \otimes \sqrt{\tfrac{1+\widehat{p}_z}{2}} \begin{pmatrix} 1 & \frac{\widehat{p}_r}{1+\widehat{p}_z} \\ \frac{\widehat{p}_l}{1+\widehat{p}_z} & -1 \end{pmatrix}, \tag{18}$$

where $\widehat{p}_z = p_z/p$, $\widehat{p}_r = (p_x - ip_y)/p$, and $\widehat{p}_l = (p_x - ip_y)/p$, we obtain the eigen solutions of the Dirac equation in the Dirac-Pauli γ-matrix representation as follows

$$\Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \widehat{u}\widetilde{\Psi}_{p\mu\nu}. \tag{19}$$

The eigen solutions of the Dirac equation are

$$\Psi_{p++}(\boldsymbol{r}, s, \tau) = \mathcal{N} \begin{pmatrix} 1 \\ \frac{\widehat{p}_l}{1+\widehat{p}_z} \\ \frac{cp}{E+mc^2} \\ \frac{\widehat{p}_l}{1+\widehat{p}_z}\frac{cp}{E+mc^2} \end{pmatrix} e^{i\boldsymbol{p}\cdot\boldsymbol{r}}, \tag{20}$$

$$\Psi_{p-+}(\boldsymbol{r}, s, \tau) = \mathcal{N} \begin{pmatrix} \frac{\widehat{p}_r}{1+\widehat{p}_z} \\ -1 \\ \frac{-\widehat{p}_r}{1+\widehat{p}_z}\frac{cp}{E+mc^2} \\ \frac{cp}{E+mc^2} \end{pmatrix} e^{i\boldsymbol{p}\cdot\boldsymbol{r}}, \tag{21}$$

$$\Psi_{p+-}(\boldsymbol{r}, s, \tau) = i\mathcal{N} \begin{pmatrix} \frac{-cp}{E+mc^2} \\ \frac{\widehat{p}_l}{1+\widehat{p}_z}\frac{cp}{E+mc^2} \\ 1 \\ \frac{\widehat{p}_l}{1+\widehat{p}_z} \end{pmatrix} e^{i\boldsymbol{p}\cdot\boldsymbol{r}}, \tag{22}$$

$$\Psi_{p--}(\boldsymbol{r}, s, \tau) = i\mathcal{N} \begin{pmatrix} \frac{\widehat{p}_r}{1+\widehat{p}_z}\frac{cp}{E+mc^2} \\ \frac{-cp}{E+mc^2} \\ \frac{\widehat{p}_r}{1+\widehat{p}_z} \\ -1 \end{pmatrix} e^{i\boldsymbol{p}\cdot\boldsymbol{r}}, \tag{23}$$

where

$$\mathcal{N} = \tfrac{1}{(2\pi\hbar)^{3/2}} \sqrt{\tfrac{E+mc^2}{2E}} \sqrt{\tfrac{1+\widehat{p}_z}{2}}. \tag{24}$$

The 4×4 helicity operator now becomes

$$\Sigma_\parallel = \mathbb{I}_\tau \otimes \begin{pmatrix} \widehat{p}_z & \widehat{p}_r \\ \widehat{p}_l & -\widehat{p}_z \end{pmatrix}, \tag{25}$$

with

$$\Sigma_\parallel \Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \mu \Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau). \tag{26}$$

The 4×4 particle-antiparticle operator now reads

$$T_3 = \frac{1}{E} \begin{pmatrix} mc^2 & 0 & cp\,\widehat{p}_z & cp\,\widehat{p}_r \\ 0 & mc^2 & cp\,\widehat{p}_l & -cp\,\widehat{p}_z \\ cp\,\widehat{p}_z & cp\,\widehat{p}_r & -mc^2 & 0 \\ cp\,\widehat{p}_l & -cp\,\widehat{p}_z & 0 & -mc^2 \end{pmatrix}, \tag{27}$$

with

$$T_3 \Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \nu \Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau). \tag{28}$$

The Hamiltonian of the Dirac equation is $\widehat{H} = ET_3$, and

$$\widehat{H}\Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \nu E\Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau). \tag{29}$$

The particle-antiparticle unitary transformation now reads

$$C = \widehat{u}\widetilde{C}\widehat{u}^{-1} = -\mathbb{I}_\sigma \otimes \tau_2, \tag{30}$$

which transforms a particle state to an antiparticle state with the helicity intact, and *vice versa*,

$$C\Psi_{p\mu\nu}(\boldsymbol{r}, s, \tau) = \Psi_{p\mu-\nu}(\boldsymbol{r}, s, \tau). \tag{31}$$

Again, the Hamiltonian $\widehat{H}$ is anti-commutative with the particle-antiparticle transformation C, $C\widehat{H}C^{-1} = -\widehat{H}$, and $\{\widehat{H}, C\} = 0$.

Dirac particles in central potentials: (A) $\widehat{\kappa}$ — the analog of the helicity in the central potential. The Hamiltonian of a Dirac particle in a central vector potential $V_v(r)$ reads

$$\widehat{H} = c\boldsymbol{\alpha} \cdot \boldsymbol{p} + mc^2\beta + V_v(r). \tag{32}$$

For a free Dirac particle, the conserved linear momentum of the particle specifies a special direction of the system. The spin alinment of the Dirac particle along the linear momentum results in the helicity which is conserved consistently with the linear momentum. The nature of the spin alignment is closely related to the linearization of the squared relativistic energy operator

$$[\widehat{H} - V_v(r)]^2 = [mc^2]^2 + (cp)^2,\ (cp)^2 = -(c\hbar)^2\nabla^2. \tag{33}$$

For the free particle, the central potential $V_v(r)$ in Eq. (33) vanishes and

$$\widehat{H}^2 = [mc^2]^2 + (cp)^2. \tag{34}$$

Since the linear momentum is conserved, the linearization should be carried out in Cartesian coordinates, and one gets the Dirac Hamiltonian Eq. (1). This means that the linearization of the squared free particle Hamiltonian operator in Cartesian coordinates leads to the helicity operator and the unique spin alignment of the free particle along its linear momentum direction naturally leads to the helicity conservation.

In a central potential, the particle has certain angular momentum, and the natural frame is the spherical coordinates. In this case, the squared kinetic energy operator reads

$$(cp)^2 = -(c\hbar)^2\nabla^2 = -(c\hbar)^2\left[\tfrac{1}{r^2}\tfrac{d}{dr}r^2\tfrac{d}{dr} - \tfrac{\widehat{L}^2}{r^2}\right]. \tag{35}$$

The problem is how to linearize $[\widehat{H} - V_v(r)]^2$ in the spherical coordinates. This has been done and the result is well known as follows,

$$\widehat{H} - V_v(r) = \beta mc^2 + \alpha_r\left(\widehat{p}_r + \tfrac{ic\hbar}{r}\beta\widehat{\kappa}\right), \tag{36}$$

$$\begin{aligned}\widehat{H} &= \alpha_r\left(c\widehat{p}_r + \tfrac{ic\hbar}{r}\beta\widehat{\kappa}\right) + V_v(r) + \beta mc^2 \\ &= \begin{pmatrix} mc^2 + V_v & c\widehat{p}_r\sigma_r - \frac{ic\hbar}{r}\sigma_r\widehat{\kappa} \\ c\widehat{p}_r\sigma_r + \frac{ic\hbar}{r}\sigma_r\widehat{\kappa} & -mc^2 + V_v \end{pmatrix},\end{aligned} \tag{37}$$

where

$$\alpha_r = \tfrac{\widehat{\alpha}\cdot\widehat{r}}{r} = \begin{pmatrix} 0 & \sigma_r \\ \sigma_r & 0 \end{pmatrix}, \sigma_r = \tfrac{\boldsymbol{\sigma}\cdot\boldsymbol{r}}{r} = \sqrt{\tfrac{3}{4\pi}}\textstyle\sum_\mu Y_{1\mu}\sigma_\mu, \tag{38}$$

with

$$\sigma_0 = \sigma_z,\ \sigma_\pm = \tfrac{1}{2}(\sigma_x \pm i\sigma y),\ \widehat{p}_r = -i\hbar(\tfrac{d}{dr} + \tfrac{1}{r}), \tag{39}$$

and $\widehat{\kappa}$ appearing in $\widehat{H}$ is just referred to its eigen values $\pm(j + \frac{1}{2})$ symbolically, it is in fact a 4×4 operator

$$\hbar\widehat{\kappa} = \beta(\boldsymbol{\Sigma} \cdot \boldsymbol{L} + \hbar) = \begin{pmatrix} \boldsymbol{\sigma} \cdot \boldsymbol{L} + \hbar & 0 \\ 0 & -(\boldsymbol{\sigma} \cdot \boldsymbol{L} + \hbar) \end{pmatrix}. \tag{40}$$

This means that $\widehat{\kappa}$ comes from the linearization of the operator $[\widehat{H} - V_v(r)]^2$ in the spherical coordinates and $\widehat{\kappa}$ is thus the analog of the helicity for the Dirac particle in the central potentials. In fact, in the central potentials, the only characteristic direction of the system is related to its angular momentum, so that the spin can only make its alignment along the angular momentum with two possibilities for a given total angular momentum j: $j = l + \frac{1}{2}$ for parallel alignment and $j = (l+1) - \frac{1}{2}$ for anti-parallel alignment. The mixing between the harmonic spin functions $\Phi^A_{jm} \equiv \sum_{m_l, m_s} C^{jm}_{lm_l 1/2 m_s} Y_{lm_l}(\Omega)\chi_{m_s}(s)$ and $\Phi^B_{jm} \equiv \sum_{m_l, m_s} C^{jm}_{l+1 m_l 1/2 m_s} Y_{l+1 m_l}(\Omega)\chi_{m_s}(s)$ is due to the scalar operator α_r contains the 1-rank spherical harmonic function $Y_{1\mu}$ and the 1-rank tensors $\sigma_\mu(\mu = 0, \pm 1)$ which change angular momentum by $\Delta l = \pm 1$ and keep the total angular momentum jm conserved.

Since in the γ-splitting scheme, the parity operator $i\gamma_4 = I_\sigma \otimes \tau_3$ indicates that the particle and antiparticle components have opposite intrinsic parities which, together with the opposite parities of the components Φ^A_{jm} and Φ^B_{jm}, make the Dirac particle in central potentials have a definite total parity.

(B) The particle-antiparticle symmetry of Dirac equation in central potentials. To study the particle-antiparticle symmetry of Dirac equation in central potentials, we add a scalar central potential $V_s(r)$ in the Dirac Hamiltonian,

$$\begin{aligned}\widehat{H} &= \alpha_r\left[c\widehat{p}_r + \tfrac{ic\hbar}{r}\beta\widehat{\kappa}\right] + V_v(r) + \beta[mc^2 + V_s(r)] \\ &= \begin{pmatrix} (mc^2 + V_s) + V_v & c\widehat{p}_r\sigma_r - \frac{ic\hbar}{r}\sigma_r\widehat{\kappa} \\ c\widehat{p}_r\sigma_r + \frac{ic\hbar}{r}\sigma_r\widehat{\kappa} & -(mc^2 + V_s) + V_v \end{pmatrix}.\end{aligned} \tag{41}$$

Let's introduce the particle-antiparticle unitary transformation

$$C = \begin{pmatrix} 0 & i\sigma_r \\ -i\sigma_r & 0 \end{pmatrix} = C^\dagger = C^{-1}, \tag{42}$$

which is just a generalization of the unitary particle-antiparticle transformation for the free Dirac particle. The transformed Hamiltonian is

$$\widetilde{H} = C\widehat{H}C^{-1} = -\widehat{H}. \tag{43}$$

Here we have used the transformation property of the scalar potential and the vector potential as usual

$$CV_sC^{-1} = V_s, \quad CV_vC^{-1} = -V_v \tag{44}$$

The Hamiltonian $\widehat{H}$ of the system is again anti-commutative with the unitary particle-antiparticle transformation C, $\{\widehat{H}, C\} = 0$.

Under the transformation C, $\widetilde{\Psi}_{njm\kappa\nu} = C\Psi_{njm\kappa\nu}$, the Dirac equations for the particle and antiparticle read,

$$\widehat{H}\Psi_{njm\kappa\nu} = E\Psi_{njm\kappa\nu}, \tag{45}$$

$$\widehat{H}\widetilde{\Psi}_{njm\kappa\nu} = -E\widetilde{\Psi}_{njm\kappa\nu}. \tag{46}$$

Explicitly,

$$\Psi_{njm++} = \begin{pmatrix} \Phi^A_{jm} f(r) \\ \Phi^B_{jm} ig(r) \end{pmatrix}, \Psi_{njm+-} = i\begin{pmatrix} \Phi^A_{jm} g(r) \\ \Phi^B_{jm} if(r) \end{pmatrix}, \tag{47}$$

$$\Psi_{njm-+} = \begin{pmatrix} \Phi^B_{jm} f(r) \\ \Phi^A_{jm} ig(r) \end{pmatrix}, \Psi_{njm--} = i\begin{pmatrix} \Phi^B_{jm} g(r) \\ \Phi^A_{jm} if(r) \end{pmatrix}. \tag{48}$$

Since the particle-antiparticle transformation exchanges the radial functions $f(r)$ and $g(r)$ between the two components of the eigen functions of the Dirac equation and keeps the total angular momentum (spherical spinors) of the eigen functions Φ^A_{jm} and Φ^B_{jm} intact, the transformed Hamiltonian leads to the same coupled eigen equations for the radial functions $f(r)$ and $g(r)$ of the antiparticle as that for the particle. Therefore the antiparticle has the same radial eigen functions as the particle does, but these two eigen radial functions exchange their position in their eigen solutions.

In the above, besides the quantum number κ as the eigen values of $\hat{\kappa}/(j+1/2)$ ($\kappa = \pm 1$ instead of $\pm(j+1/2)$), we have introduced the particle-antiparticle quantum numbers ν ($= \pm 1$). The eigen function of the Dirac particles in the scalar and vector central potentials have been written as $\Psi_{njm\kappa\nu}$ with the following dual symmetry and the conserved quantum numbers,

$$\begin{aligned}
\widehat{C}\Psi_{njm\kappa\nu} &= \widetilde{\Psi}_{njm\kappa\nu} = \Psi_{njm\kappa-\nu}, && (49)\\
\widehat{H}\Psi_{njm\kappa\nu} &= \nu E_{nj\kappa}\Psi_{njm\kappa\nu}, && (50)\\
\widehat{J}^2\Psi_{njm\kappa\nu} &= j(j+1)\Psi_{njm\kappa\nu}, && (51)\\
\widehat{J}_z\Psi_{njm\kappa\nu} &= m\Psi_{njm\kappa\nu}, && (52)\\
\widehat{\kappa}/(j+1/2)\Psi_{njm\kappa\nu} &= \kappa\Psi_{njm\kappa\nu}. && (53)
\end{aligned}$$

It is remarkable that the anti-symmetry of a system does not lead to a conserved operator because the anti-symmetry operator is not commutative but anti-commutative with the Hamiltonian. However, as the particle-antiparticle space is introduced, the discrete group $(\mathbb{I}, C)$ can be embedded in the enlarged continuous intrinsic particle-antiparticle space as a discrete subgroup of the enlarged continuous group with the generators (τ_i) and one can find the invariant operators and the corresponding quantum numbers. For example, in the cases investigated we do find both the anti-symmetry operator C and the conserved particle-antiparticle operator T_3.

Summary. We find that the internal γ-degrees of freedom in relativistic quantum mechanics and quantum field theory is a composite and coherent combination of the spin degrees of freedom and the particle-antiparticle degrees of freedom. This coherent combination can only be realized for free particles and it hides the particle-antiparticle degrees of freedom. As the coherent composite γ-degrees of freedom are split by interactions, the γ space is broken into the spin σ-space and particle-antiparticle τ-space. In general cases with interactions, one needs to describe the Dirac particles in the σ-space and τ-space separately, which provides an alternative approach besides the nonlinear Clifford algebra or Dirac ring. This implies that the splitting of the coherent composite γ-space into its constituent σ-space and τ-space is just the reflection of the breaking of its coherence by interactions. The advantage of the splitting is that it is more "microscopic" than Dirac ring and it leads to new and more detailed information of the wave function and the corresponding quantum number because it introduces a new intrinsic space—the particle-antiparticle space. For example, for the ortho-positronium, quite different from the conventional description, one can write down explicitly its more detailed wave function with the quantum numbers in configuration space, spin space, and particle-antiparticle space as $(l = 0, s = 1, \tau = 0)$, thus C-parity is negative (since the τ wave function is anti-symmetric); while for the para-positronium, the quantum numbers are $(l = 0, s = 0, \tau = 1)$ and C-parity is positive (since the τ wave function is symmetric). Thus explanation of the decay properties of positronium is more microscopic and transparent than the conventional theory [4].

S.J.W. and S.G.Z. are indebted to Max-Planck Institute for Nuclear Physics and particularly to Professor Hans-Christian Pauli for their hospitality. S.J.W. was partly supported by the National Natural Science Foundation of China under Grant Nos. 10375039 and 10175029. S.G.Z. was partly supported by the National Natural Science Foundation of China under Grant No. 10475003, the Major State Basic Research Development Program of China under contract No. G20000774 and the Knowledge Innovation Project of Chinese Academy of Sciences under contract No. KJCX2-SW-N02.

---

[*] Electronic address: sgzhou@itp.ac.cn; URL: `http://www.itp.ac.cn/~sgzhou`

[1] J.D. Bjørken and S.D. Drell, *Relativistic Quantum Mechanics* and *Relativistic Quantum Fields*, McGraw-Hill Book Company, New York, 1964 and 1965.

[2] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill Book Company, New York, 1980.

[3] P. Roman, *Theory of Elementary Particles*, North-Holland Publishing Company, Amsterdam, 1960.

[4] T.D. Lee, *Particle Physics and Introduction to Field Theory*, Harwood Academic Publishers, 1983.